



**IN THE MATTER OF** an Application  
for a French Patent  
in the name of  
**CENTRE NATIONAL DE LA RECHERCHE  
SCIENTIFIQUE (CNRS)**  
filed under No. 99 08772, and

I, Nicolas Torno,  
c/o Cabinet REGIMBEAU, 20, Rue de Chazelles, 75847 PARIS, France,  
do solemnly and sincerely declare that I am conversant with the French and English languages  
and I am a competent translator thereof, and that the following is, to the best of my knowledge  
and belief, a true and correct translation of the Patent Application filed under No. 99 08772

by CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE (CNRS)

in FRANCE on 7 July 1999

for "Method for producing Oligosaccharides"

and the Official Certificate attached thereto.

Date: 25 June 2008

  
N. TORNO

## METHOD FOR PRODUCING OLIGOSACCHARIDES

The present invention relates to the microbiological  
5 production of oligosaccharides of biological interest.

It is now well-established that oligosaccharides play  
an important biological role especially as regards the  
activity and function of proteins; thus, they serve to  
10 modulate the half-life of proteins, and occasionally  
they are involved in the structure of the protein.  
Oligosaccharides play an essential role in antigen  
variability (for example blood groups), and in certain  
bacterial infections such as those caused by *Neisseria*  
15 *meningitidis*.

As oligosaccharides are usually obtained in a low yield  
by purification starting from natural sources, the  
synthesis of oligosaccharides has become a major  
20 challenge of carbohydrate chemistry, so as to supply  
sufficient amounts of well-characterized oligosac-  
charides, required for fundamental research or for any  
other potential applications (Boons et al., 1996).

25 The synthesis of complex oligosaccharides of biological  
interest may be performed chemically, enzymatically or  
microbiologically.

Despite the development of new chemical methods for  
30 synthesizing oligosaccharides in the course of the last  
20 years, the chemical synthesis of oligosaccharides  
remains very difficult on account of the numerous  
selective protection and deprotection steps, the  
lability of the glycoside linkages, the difficulties in  
35 obtaining regiospecific couplings, and the low  
production yields. As the number of steps increases  
with the size of the oligosaccharide, the preparation  
of large quantities of oligosaccharides longer than  
trisaccharides is no simple matter. Contrary to the

experience of peptide synthesis or nucleic acid synthesis, traditional synthetic organic chemistry cannot at the present time provide a high-quality and large-quantity synthesis of oligosaccharides, even of  
5 simple formula.

Consequently, the enzymatic methods have become more popular since they allow a regioselective synthesis under mild conditions and without a step for protection  
10 of the hydroxyl groups. The development of the enzymatic approach was made possible by the cloning and functional identification of numerous genes encoding the enzymes involved in the synthetic pathway of oligosaccharides. Thus, various types of enzyme may be  
15 used for the *in vitro* synthesis of oligosaccharides. The physiological function of the glycosyl-hydrosylases and of the glycosyl-phosphorylases is to depolymerize the oligosaccharides, but they may also be used *in vitro* in the synthesis of oligosaccharides by  
20 controlling the reaction equilibrium and kinetics. The substrates of the enzymes for these reactions are readily available, but these enzymatic reactions are not very versatile. Another enzymatic method developed uses the glycosyl-transferases of the Leloir  
25 biochemical pathway, which show strong regiospecificity for the precursor and also for the donor substrate; these glycosyl-transferases are not as readily available as the glycosyl-hydrolases. The recombinant DNA technique has recently made it possible to clone  
30 and produce a certain number of them. However, the main limitation of this enzymatic method lies in the very high cost of the sugar-nucleotides that are the sugar donors used by these enzymes.

35 The microbiological route for producing recombinant oligosaccharides *in vivo* is the most appealing of the synthetic routes since the bacterium is simultaneously responsible for the biosynthesis of the enzymes, the

regeneration of the sugar-nucleotides and, finally, the production of the oligosaccharide.

5 The first descriptions of the microbiological synthesis of oligosaccharides using recombinant bacteria may be considered to a certain extent as the studies which led to the elucidation of the pathways for the biosynthesis of the nodulation factors; these factors are signal molecules secreted by the rhizobia to allow recognition  
10 by leguminous plants in the nodulation process. Nodulation factors consist of a chito-oligosaccharide backbone bearing various substitutions. The functional identification of the *nod* genes involved in the biosynthesis of the nodulation factors was partly  
15 performed by identifying the oligosaccharides formed *in vivo* in strains of *Escherichia coli* expressing these various *nod* genes (Gérémia *et al.*, 1994; Kamst *et al.*, 1995; Spaink *et al.*, 1994; Mergaert *et al.*, 1995). However, the production of oligosaccharides per se was  
20 not the aim of these studies; these products were synthesized only in trace amounts and were identified only by means of using radioactive precursors.

On the other hand, it was recently demonstrated in our  
25 laboratory (Samain *et al.*, 1997) that the culturing at high cell density of *Escherichia coli* strains containing the *nodC* (chito-oligosaccharide synthase) gene made it possible to produce large amounts, of greater than 2 g/l, of "recombinant" chito-oligosac-  
30 charides.

However, this technique of microbiological synthesis of oligosaccharides remains limited to the production only of chito-oligosaccharides, due to the unique property  
35 of *nodC* (chito-oligosaccharide synthase) of functioning without a precursor; specifically, the other enzymes glycolyze a specific precursor and their activity is thus dependent on the presence of this precursor in the cell. The problem of the precursor is thus the main

obstacle blocking the development of the method and its extension to the production of other types of oligosaccharide.

5 One subject of the present invention is thus a method for producing an oligosaccharide of interest by a cell starting with at least one exogenous precursor internalized by said cell, said precursor being involved in the biosynthetic pathway of said  
10 oligosaccharide, said method comprising the steps (i) of obtaining a cell that comprises at least one gene encoding an enzyme capable of modifying said exogenous precursor or one of the intermediates in the biosynthetic pathway of said oligosaccharide from said  
15 exogenous precursor necessary for the synthesis of said oligosaccharide from said precursor, and also the components for expressing said gene in said cell, said cell lacking any enzymatic activity liable to degrade said oligosaccharide, said precursor and said  
20 intermediates; (ii) of culturing said cell in the presence of at least one said exogenous precursor, under conditions enabling the internalization of said exogenous precursor by said cell and the production of said oligosaccharide by said cell.

25 According to one particular embodiment, the present invention relates to a method as described above, characterized in that said cell also comprises at least one gene encoding an enzyme capable of modifying an  
30 endogenous precursor involved in the biosynthetic pathway of said oligosaccharide, said enzyme being identical to or different than the enzyme used in the method described above, and also to the components for expressing said gene in said cell and characterized in  
35 that said cell lacks any enzymatic activity liable to degrade said precursor.

The term "oligosaccharides" is intended to denote linear or branched polymers with a variable number of

residues, linkages and subunits; the number of residues being greater than 1. Oligosaccharides are carbohydrates that become converted on hydrolysis into several monosaccharide molecules; the monosaccharides  
5 being sugars that cannot be converted into a simpler substance by hydrolysis. Monosaccharides are subdivided into trioses, tetroses, pentoses, hexoses and heptoses depending on the number of carbon atoms in their hydrocarbon-based chain, and also into aldoses and  
10 ketoses depending on the presence of an aldehyde function or a ketone function in their molecule. Among the monosaccharides that are most frequently encountered, mention may be made of mannose, glucose, galactose, N-acetylglucosamine and N-acetyl-  
15 galactosamine. The number of chains of stereoisomeric oligosaccharides is extremely large, due to the large number of asymmetric carbons in the hydrocarbon-based chain.

20 The expression "exogenous precursor" is intended to denote a compound involved in the biosynthetic pathway of the oligosaccharide according to the invention that is internalized by said cell. The expression "endogenous precursor" is intended to denote a compound  
25 involved in the biosynthetic pathway of the oligosaccharide according to the invention that is naturally present in said cell.

The method according to the invention is characterized  
30 in that said cell is a cell chosen from bacteria and yeasts. According to one preferred embodiment of the invention, the bacterium is chosen from the group composed of *Escherichia coli*, *Bacillus subtilis*, *Campylobacter pylori*, *Helicobacter pylori*,  
35 *Agrobacterium tumefaciens*, *Staphylococcus aureus*, *Thermophilus aquaticus*, *Azorhizobium caulinodans*, *Rhizobium leguminosarum*, *Neisseria gonorrhoeae* and *Neisseria meningitis*. According to one preferred embodiment of the invention, the bacterium is

*Escherichia coli*. According to another embodiment of the invention, the cell is a yeast that is preferably *Saccharomyces cerevsae*, *Saccharomyces pombe* or *Candida albicans*. The cell according to the invention lacks any  
5 enzymatic activity liable to degrade said oligosaccharide, said precursor or said metabolic intermediates.

The nucleic acid sequence encoding the enzyme according  
10 to the invention is either naturally present in said cell or is introduced into said cell by the recombinant DNA techniques known to those skilled in the art. In the present description, the term "nucleic acid" will be intended to denote a DNA fragment, which is either  
15 double-stranded or single-stranded, or products of transcription of said DNAs, and/or an RNA fragment. According to one preferred embodiment, the nucleic acid sequence which is introduced into said cell by the recombinant DNA techniques and which encode an enzyme  
20 involved in the biosynthetic pathway of the oligosaccharide of interest is heterologous. The expression "heterologous nucleic acid sequence" is intended to denote a nucleic acid sequence that is not naturally present in the cell according to the  
25 invention. The heterologous nucleic acid sequence according to the invention may originate from any animal or plant, eukaryotic or prokaryotic cell type and may originate from viruses.

30 Among the prokaryotic cells from which the heterologous nucleic acid sequence originates, mention should be made of bacteria and in particular *Escherichia coli*, *Bacillus subtilis*, *Campylobacter pylori*, *Helicobacter pylori*, *Agrobacterium tumefaciens*, *Staphylococcus aureus*,  
35 *Thermophilus aquaticus*, *Azorhizobium caulinodans*, *Rhizobium leguminosarum*, *Rhizobium meliloti*, *Neisseria gonorrhoeae* and *Neisseria meningitis*.

Among the unicellular eukaryotic cells from which the heterologous nucleic acid sequence originates, mention should be made of yeasts and in particular *Saccharomyces cerevisiae*, *Saccharomyces pombe* and  
5 *Candida albicans*.

According to one preferred embodiment, the heterologous nucleic acid sequence originates from plant or animal eukaryotic cells. According to an even more preferred  
10 embodiment, the heterologous nucleic acid sequence originates from mammalian cells and preferably from human cells.

According to one preferred embodiment of the invention,  
15 the cell according to the invention is the bacterium *Escherichia coli* and the nucleic acid sequence introduced into the bacterium and encoding the enzyme according to the invention preferably originates from a bacterium chosen from the group mentioned above.

20 According to one preferred embodiment of the invention, the nucleic acid sequence encoding the enzyme according to the invention is introduced into said cell in the form of an expression vector. The vector must comprise  
25 a promoter, translation start and stop signals, and also regions suitable for regulating transcription. The vector must be able to be maintained stably in the cell over successive generations and can optionally contain particular signals specifying the secretion of the  
30 translated enzyme. These various control signals are chosen as a function of the host cell used. To this end, the nucleic acid sequences may be inserted into autonomous replication vectors within the chosen host or into integrative vectors which become integrated  
35 into the genome of the chosen host. Such vectors are prepared according to the methods commonly used by those skilled in the art, and the clones resulting therefrom may be introduced into a suitable host cell



by standard methods such as, for example, heat shock or electroporation.

5 The invention is also directed toward the above cells, characterized in that they are transformed by at least one recombinant isolated nucleic acid encoding the enzyme according to the invention or by at least one recombinant vector as defined above.

10 The method according to the invention is characterized in that said modification made by said enzyme is chosen from glycosylation, sulfatation, acetylation, phosphorylation, succinylation, methylation and addition of an enolpyruvate group. More particularly,  
15 the method according to the invention is characterized in that said enzyme is an enzyme capable of carrying out a glycosylation, which is chosen from glycosyl-transferases, glycosyl-hydrolases and glycosyl-phosphorylases. According to one preferred embodiment,  
20 the enzyme capable of carrying out the glycosylation is a glycosyl-transferase. According to one preferred embodiment, the glycosyl-transferase according to the invention is chosen from  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase,  $\beta$ -1,4-galactosyl-transferase,  $\alpha$ -1,3-galactosyl-transferase,  $\alpha$ -1,4-galactosyl-transferase,  $\alpha$ -2,3-sialyl-transf $\acute{e}$ rase, 1'  $\alpha$ -1,3-fucosyl-transf $\acute{e}$ rase.  
25 The glycosyl-transferases used in the present invention are capable of stereospecific conjugation of specific activated saccharide units on a specific acceptor  
30 molecule. The activated saccharides generally consist of uridine diphosphate, guanosine diphosphate and cytidine diphosphate saccharide derivatives. Thus, the activated saccharides may be a UDP-saccharide, a GDP-saccharide or a CMP-saccharide.

35 Certain genes encoding glycosyl-transferases used in the method according to the invention have been described previously; thus, international patent application WO 96/10086 describes the standard

oligosaccharide synthesis: in a first step, the various glycosyl-transferases are produced in recombinant bacteria containing the *lgtA*, *lgtB* and *lgtC* genes of *Neisseria gonorrhoeae*, and, after purifying the recombinant enzymes thus produced, the oligosaccharides are synthesized *in vitro* in the presence of the required precursors and sugar-nucleotides.

According to certain embodiments of the invention, the enzyme capable of performing an acetylation is encoded by the *NodL* gene of the bacterium *Azorhizobium caulinodans*. According to another embodiment, the enzyme capable of performing a sulfatation is encoded by the *NodH* gene of the bacterium *Rhizobium meliloti*.

The method according to the invention is characterized in that said cell culturing is preferably performed on a carbon-based substrate; according to one particular embodiment of the invention, said carbon-based substrate is chosen from glycerol and glucose. Other carbon-based substrates may also be used; mention should be made of maltose, starch, cellulose, pectin and chitin. According to another embodiment, the cell culturing is performed on a substrate composed of amino acids and/or protein and/or lipids.

The method according to the invention is characterized in that said culturing step is performed under conditions allowing the production of a culture with a high cell density; this culturing step comprises a first phase of exponential cell growth ensured by said carbon-based substrate, a second phase of cell growth limited by said carbon-based substrate which is added continuously, and finally a third phase of slowed cell growth obtained by continuously adding to the culture an amount of said substrate that is less than the amount of substrate added in step b) so as to increase the content of oligosaccharides produced in the high cell density culture.

The method according to the invention is characterized in that the amount of substrate added continuously to the cell culture during said phase c) is at least 30% less, preferentially 50% and preferably 60% less than the amount of substrate added continuously during said phase b). The method according to the invention is also characterized in that said exogenous precursor is added during phase b).

According to one embodiment of the invention, the method is characterized in that said exogenous precursor is of carbohydrate nature, preferably of oligosaccharide nature.

The novelty and feasibility of the method according to the invention is based on the use of two modes of internalization of the exogenous precursor that do not destroy the integrity of the cell or attack its vital functions. This especially excludes the standard techniques of membrane permeabilization with organic solvents which block growth and energy metabolism. The two possible modes for internalizing the exogenous precursor use a passive or active transport mechanism.

The invention relates firstly to a method that is characterized in that said exogenous precursor is internalized according to a passive transport mechanism. The expression "internalization by passive transport" is intended to denote the passive diffusion of any of the exogenous precursor across the plasma membrane, the molecular flow being oriented from the zones of highest concentration to the zones of lowest concentration so as to tend finally toward a state of equilibrium. The internalization by passive transport consists in using an exogenous precursor that is small enough and hydrophobic enough to diffuse passively across the membrane. A monosaccharide precursor whose anomeric position is blocked with an alkyl substitute

constitutes an example of a precursor that may be internalized in this manner. The present invention thus relates to a method that is characterized in that said exogenous precursor is a monosaccharide whose anomeric carbon is linked to an alkyl group; preferably, said alkyl group is an allyl group. One of the objects of the invention is thus to provide a method for producing oligosaccharides that contain a functionalizable group such as the allyl group and that can consequently be used as precursors for the chemical synthesis of glycoconjugates (neoglycoprotein or neoglycolipids) or glycopolymers. The reason for this is that the double bond of the allyl group is able to be opened by ozonolysis to form an aldehyde and to allow the oligosaccharide to conjugate onto a protein by reductive amination (Roy *et al.*, 1997). Another route is the addition of cysteamine (Lee and Lee, 1974, Roy *et al.*, 1997) to the allylic double bond to form an amine end group which may react, for example, with the carboxylic groups of proteins.

According to one particular embodiment, the method according to the invention concerns the production of ( $\beta$ -D-Gal-[1 $\rightarrow$ 4]- $\beta$ -D-GlcNac-1 $\rightarrow$ O-allyl); the method is characterized in that said cell is a bacterium of *LacZ*<sup>-</sup> genotype, said enzyme is  $\beta$ -1,4-galactosyl-transferase, said substrate is glycerol and said precursor is allyl-N-acetyl- $\beta$ -D-glucosaminide ( $\beta$ -D-GlcNac-1 $\rightarrow$ O-allyl). Finally, according to another particular embodiment, the method according to the invention is characterized in that the double bond of the allyl group of said ( $\beta$ -D-Gal-[1 $\rightarrow$ 4]- $\beta$ -D-GlcNac-1 $\rightarrow$ O-allyl) is chemically modified by addition, oxidation or ozonolysis reactions.

The present invention also relates to a method that is characterized in that said precursor is internalized according to an active transport mechanism. The expression "internalization by active transport" is

intended to denote the ability of cells and preferably of bacteria to selectively admit and concentrate certain exogenous substances or precursors into their cytoplasm. This transport is performed by transporters of protein nature known as permeases, which act as enzymes; permeases are inducible catalysts, that is to say catalysts that are synthesized in the presence of the substrate or the precursor. According to one particular embodiment of the invention, lactose and  $\beta$ -galactosides constitute precursors that are actively transported into the cytoplasm of the bacterium *Escherichia coli* by lactose permease, also known as galactoside permease. The invention thus relates to a method according to the invention that is characterized in that said active transport of said precursor is performed by lactose permease. Lactose permease has fairly broad specificity, which allows it to transport molecules other than lactose.

The reason for this is that it is capable of transporting various natural or synthetic  $\beta$ -galactosides,  $\alpha$ -galactosides and sucrose. One of the objects of the invention is thus to provide, according to a preferred embodiment, a method that is characterized in that said precursor is lactose, which constitutes the base moiety for a great many biologically active oligosaccharides. It is also within the scope of the invention to provide a method that is characterized in that said precursor is chosen from the group composed of: (i) natural or synthetic  $\beta$ -galactosides, preferably from 4-O- $\beta$ -D-galactopyranosyl-D-fructofuranose (lactulose), 3-O- $\beta$ -D-galactopyranosyl-D-arabinose and allyl- $\beta$ -D-galactopyranoside, (ii)  $\alpha$ -galactosides, preferably melibiose and raffinose, (iii) sucrose.

The specificity of lactose permease may even be modified by mutation and allow the transport of other compounds such as maltose and cellobiose. All these

compounds may thus be used as precursors for the synthesis of oligosaccharides. It is also within the scope of the invention to use as precursors lactose analogs containing a chemically reactive group for a subsequent functionalization of the product; preferably, one of these analogs is allyl- $\beta$ -D-galactopyranoside. It is also within the scope of this invention to use other permeases possibly modified by recombinant DNA techniques to allow the internalization of different types of precursor.

The  $\beta$ -galactosides are normally hydrolyzed in the cytoplasm of the bacterium by the  $\beta$ -galactosidase encoded by the *LacZ* gene. In order to overcome this problem, a *lacZ*<sup>-</sup> bacterial mutant lacking  $\beta$ -galactosidase activity is used when the precursor used is lactose and/or a  $\beta$ -galactoside. One of the objects of the invention is thus also to provide the method according to the invention that is characterized in that said cell lacks enzymatic activity liable to degrade said precursor or said metabolic intermediates.

In the method according to the invention, said cell may be lacking in enzymatic activity liable to degrade said precursor(s).

According to one preferred embodiment, the method is characterized in that said cell has a genotype chosen from *LacZ*<sup>-</sup>.

According to another aspect of the invention, the method is characterized in that it also comprises the addition of an inducer to said culture medium to induce the expression in said cell of said enzyme and/or of a protein involved in said active transport; according to one preferred embodiment, the method according to the invention is characterized in that said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG) and said protein is lactose permease.

The invention makes it possible for the first time to produce complex oligosaccharides in yields of the order of one gram per liter. Depending on its size, the oligosaccharide either accumulates in the bacterial cytoplasm or is secreted into the culture medium. Thus, according to one preferred embodiment, the method according to the invention is used for the production of the trisaccharide 4-O-[3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranosyl]-D-glucopyranose, ( $\beta$ -D-GlcNac-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-[1 $\rightarrow$ 4]-D-Glc); it is characterized in that said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype, said enzyme is  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase, said substrate is glycerol, said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG) and said precursor is lactose.

According to a second preferred embodiment, the method according to the invention is used for the production of lacto-N-neo-tetraose and polylactosamine; it is characterized in that said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype, said enzymes are  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase and  $\beta$ -1,4-galactosyl-transferase, said substrate is glucose, said inducer is isopropyl- $\beta$ -D-thiogalactoside (IPTG) and said precursor is lactose.

According to a third preferred embodiment, the method according to the invention is used for the production of allyl 3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranoside, ( $\beta$ -D-GlcNac-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-1 $\rightarrow$ O-allyl); it is characterized in that said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype, said enzyme is  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase, said substrate is glycerol, said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG) and said precursor is allyl- $\beta$ -D-galactopyranoside.

According to a fourth preferred embodiment, the method according to the invention is used for the production of analogs of lacto-N-neo-tetraose and of polylactosamines in which the glucose residue is replaced with an allyl group; it is characterized in that said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype, said enzymes are  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase and  $\beta$ -1,4-galactosyl-transferase, said substrate is glucose, said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG) and said precursor is allyl- $\beta$ -D-galactopyranoside.

According to a fifth preferred embodiment, the method according to the invention is used for the production of allyl- $\beta$ -D-lactosamine ( $\beta$ -D-Gal-[1 $\rightarrow$ 4]- $\beta$ -D-GlcNac-1 $\rightarrow$ O-allyl); it is characterized in that said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype, said enzyme is  $\beta$ -1,4-galactosyl-transferase, said substrate is glycerol and said precursor is allyl-N-acetyl  $\beta$ -D-glucosaminide ( $\beta$ -D-GlcNac-[1 $\rightarrow$ O-allyl]).

The invention also relates to a method that makes it possible to envisage the production of a large number of different oligosaccharides obtained by glycosylation of lactose. Specifically, besides the genes *lgtA* and *lgtB* which respectively encode  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase and  $\beta$ -1,4-galactosyl-transferase, several genes of bacterial glycosyl-transferases using lactose as precursor have recently been cloned. These are *lgtC* ( $\beta$ -1,4-galactosyl-transferase) and *Lst* ( $\alpha$ -2,3-sialyl-transferase) (Gilbert *et al.*, 1997). Using these genes in a method according to the invention makes it possible to produce molecules such as globotriose (P<sup>k</sup> blood antigen) and sialyl-lactose. Moreover coexpression of the *lgtA* and *lgtB* genes with the gene for  $\alpha$ -1,3-fucosyl-transferase from *Helicobacter pylori* (Martin *et al.*, 1997) according to a method according to the invention makes it possible to obtain Lewis<sup>x</sup> pentasaccharide. The



addition of the *Lst* ( $\alpha$ -2,3-sialyl-transferase) gene gives access to the sialyl Lewis<sup>x</sup> hexasaccharide.

5 The method according to the invention also makes it possible to obtain a large number of different oligosaccharides obtained by glycosylation of exogenous precursors other than lactose and transported by lactose permease or by other permeases.

10 The method according to the invention makes it possible to obtain a large number of different oligosaccharides obtained by *in vivo* modification (sulfatation, acetylation, phosphorylation, succinylation, methylation, addition of an enolpyruvate group) of  
15 precursors. The synthesis of certain oligosaccharides may necessitate the modification of endogenous precursors, in addition to the modification of exogenous precursors. Thus, it may be envisaged to introduce into a K12 *Escherichia coli* bacterium the  
20 gene for the enzyme involved in the metabolism of an endogenous precursor to allow the production of certain sugar-nucleotides such as, for example, CMP-sialic acid, UDP-GalNAc or GDP-fucose, which are not normally produced by this bacterial strain, so as to achieve the  
25 synthesis of an oligosaccharide of interest. For example, UDP-GalNAc may be produced from UDP-GlcNAc if the epimerase gene is introduced into a cell according to the invention.

30 Contrary to the enzymatic method for the *in vitro* synthesis of oligosaccharides, which requires the use of very expensive molecules such as ATP, acetyl-CoA, PAPS (adenosine 3'-phosphate 5'-phosphosulfate) or phospho-enolpyruvate, one of the advantages of the  
35 present invention lies in the fact that these molecules are naturally recycled into the cell, thus making it possible to reduce the production costs of the oligosaccharides.

Another object of the invention is to provide a method for producing oligosaccharides that are labeled with or enriched in radioisotopes; such oligosaccharides are  
5 extremely precious for fundamental biological or conformational analysis studies. The invention thus relates to a method for producing an oligosaccharide that is labeled with at least one radioisotope, characterized in that said cell is cultured on said  
10 carbon-based substrate labeled with said radioisotope and/or in the presence of a said precursor labeled with said radioisotope. The radioisotopes are preferably chosen from the group composed of:  $^{14}\text{C}$ ,  $^{13}\text{C}$ ,  $^3\text{H}$ ,  $^{35}\text{S}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ .

15 The invention also relates to an oligosaccharide which may be obtained by a method according to the invention.

According to one particular embodiment, the invention  
20 relates to an activated oligosaccharide that may be used for the chemical synthesis of glycoconjugates or glycopolymers that may be obtained by a method as described above, said oligosaccharide being characterized in that the double bond of the allyl  
25 group is chemically modified by addition, oxidation or ozonolysis reactions.

The oligosaccharide according to the invention is useful in a wide range of therapeutic and diagnostic  
30 applications; it may be used, for example, as an agent for blocking cell surface receptors in the treatment of a host of diseases involving cellular adhesion, or may be used as nutritional supplements, antibacterial agents, anti-metastatic agents and anti-inflammatory  
35 agents. The invention thus relates to an oligosaccharide according to the invention as a medicinal product, and especially as a medicinal product intended for selectively preventing the adhesion of biological molecules. The oligosaccharide

according to the invention is also used as a medicinal product intended for treating cancer, inflammation, heart diseases, diabetes, bacterial infections, viral infections and neurological diseases and as a medicinal  
5 product intended for grafts. The invention also relates to a pharmaceutical composition, characterized in that it comprises an oligosaccharide according to the invention and a pharmaceutically acceptable vehicle.

10 Finally, the invention also relates to the agricultural and agronomic use of an oligosaccharide according to the invention, especially for the growth and defense of plants. Specifically, oligosaccharides play a predominant role in Rhizobium/leguminous plant  
15 symbiosis. Indeed, certain oligosaccharides originating from the hydrolysis of fungal or plant glycoproteins or walls can act as plant hormones or as elicitors of defense reactions in plants.

20 The industrial advantage of the method according to the invention is obvious since it makes it possible for the first time to achieve a production of the order of a kilogram of complex oligosaccharides of biological interest. All the oligosaccharides of biological  
25 interest that we envisage synthesizing at the industrial scale are currently available only at the mg scale and at extremely high costs (up to 1 million FFr per gram); the cost price of these compounds produced by the present microbiological route are infinitely  
30 lower.

The characteristics and advantages of the present invention will be demonstrated more clearly on reading the examples and figures which follow, the keys to  
35 which are represented below.

## FIGURES

**Figure 1: Principle of the method for producing the trisaccharide 4-O-[3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranosyl]-D-glucopyranose, ( $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-[1 $\rightarrow$ 4]-D-Glc)**

5

Lactose ( $\beta$ -D-Gal-[1-4]- $\beta$ -D-Glc) is transported into the cell by lactose permease (Lac permease). The lactose cannot be hydrolyzed in the cell since the strain is a *LacZ*<sup>-</sup> mutant. Expression of the *lgtA* gene allows the production of the LgtA enzyme which transfers a GlcNAc from UDP-GlcNAc onto a lactose molecule. The trisaccharide formed ( $\beta$ -D-GlcNAc-[1-3]- $\beta$ -D-Gal-[1-4]-D-Glc) is excreted into the medium.

10

**Figure 2: High cell density culturing of the control strain JM 109 and of the strain JM 109 (pCWlgtA) containing the glycosyl transferase gene LgtA**

25

Lactose is added continuously and the residual lactose is determined enzymatically. The concentration of hydrolyzable GlcNAc in the culture medium is measured colorimetrically after acid hydrolysis. The added lactose represents the total cumulative amount of lactose which was added continuously.

20

**Figure 3: Mass spectrum in FAB<sup>+</sup> mode of the trisaccharide 4-O-[3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranosyl]-D-glucopyranose, ( $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-[1 $\rightarrow$ 4]-D-Glc) purified from the culture supernatant of the strain JM 109 (*lgtA*)**

30

The two quasi-molecular ions [M+H]<sup>+</sup> and [M+Na]<sup>+</sup> are observed at m/z 546 and 568. An ion [M+H]<sup>+</sup> at m/z 442 is also observed, which is due to the presence of  $\beta$ -D-GlcNAc-[1-3]-IPTG. This indicates that the IPTG (isopropyl  $\beta$ -D-thiogalactose) used to induce Lac permease and LgtA is also glycosylated.

35

**Figure 4: Proton NMR spectrum of the trisaccharide 4-O-[3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranosyl]-D-glucopyranose, ( $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-[1 $\rightarrow$ 4]-D-Glc) at 323° K**

5

The signal at 1.4 ppm is due to the protons of the isopropyl group of the glycosylated IPTG derivative.

**Figure 5:  $^{13}\text{C}$  NMR spectrum of the trisaccharide 4-O-[3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranosyl]-D-glucopyranose, ( $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-[1 $\rightarrow$ 4]-D-Glc)**

10

**Figure 6: Principle of the method for producing lacto-N-neo-tetraose ( $\beta$ -D-Gal-[1-4]- $\beta$ -D-GlcNAc-[1-3]- $\beta$ -D-Gal-[1-4]- $\beta$ -D-Glc)**

15

Lactose ( $\beta$ -D-Gal-[1-4]- $\beta$ -D-Glc) is transported into the cell by Lac permease. The lactose cannot be hydrolyzed in the cell since the strain is a *LacZ<sup>-</sup>* mutant. Expression of the *lgtA* gene allows the production of the LgtA enzyme which transfers a GlcNAc from UDP-GlcNAc onto a lactose molecule. The trisaccharide formed is then used as a precursor by LgtB which transfers a galactose molecule from UDP-Gal to form lacto-N-neo-tetraose ( $\beta$ -D-Gal-[1-4]- $\beta$ -D-GlcNAc-[1-3]- $\beta$ -D-Gal-[1-4]- $\beta$ -D-Glc).

20

25

**Figure 7: High cell density culturing of the strain JM 109 (pCWlgtA, pBBlgtB)**

30

Culturing in the presence of lactose at high concentration (5 g.l<sup>-1</sup>) and at low concentration (1 g.l<sup>-1</sup>).

35

**Figure 8: Separation on Biogel P4 of the oligosaccharides produced by the strain JM 109 (pCWlgtA, pBBlgtB) in the presence of lactose at an initial concentration of 5 g.l<sup>-1</sup> (A) or of 1 g.l<sup>-1</sup> (B)**

The peaks 1, 2, 3 and 4 correspond, respectively, to lacto-N-neo-tetraose, lacto-N-neo-hexaose, lacto-N-neo-octaose and lacto-N-neo-decaose.

5

## EXAMPLES

### Example 1: Materials and methods

#### 10 1.1. Origin of the plasmids and bacterial strains

The strain JM 109 of *Escherichia coli* K12 (Yannisch-Perron et al., 1984) was used as host cells for all the oligosaccharide production examples described. The strain was obtained from the DSM (Deutsche Sammlung von Mikroorganismen). The genotype of the strain is as follows: *F traD36 lacI<sup>g</sup> Δ(lacZ)M15 proA<sup>+</sup>B<sup>+</sup>/e14<sup>-</sup>(McrA<sup>-</sup>) Δ(lac-proAB) supE44 recA1 endA1 gyrA96 (Nal<sup>r</sup>) thi hsdR17 relA1*.

20

The *lgtA* and *lgtB* genes of *Neisseria meningitis* MC58 were supplied by Dr W. Wakarchuk (Institute for Biological Sciences, National Research Council of Canada, 100 Sussex Drive, Ottawa, Ontario, K1A 0R6, Canada) in the form of two plasmids pCW, one containing the *lgtA* gene (referred to herein as pCWlgtA) and the other containing the *lgtB* gene (referred to herein as pCWlgtB). The sequences of these two genes are available from the GenBank databank under the number U25839. The plasmid pLitmus28 was purchased from the company New England's Biolabs. The plasmid pBBR1MCS was supplied by Dr M. Kovach (Department of Microbiology and Immunology, Louisiana State University, Shreveport, LA 71130-3932, USA).

35

#### 1.2. Subclonings

We used the standard techniques of molecular biology described by Sambrook et al. (1989).

The 0.835-kb DNA fragment containing the *lgtB* gene was obtained by digestion of the plasmid pCWLgtB with *Bam*HI and *Hind*III. This fragment was subcloned into the vector pLitmus28 predigested with *Bam*HI and *Hind*III to form the plasmid pLitlgtB. The 0.9-kb fragment containing the *lgtB* gene was excised from the plasmid pLitlgtB by digestion with *Xho*I and *Hind*III and subcloned into the plasmid pBBR1MCS predigested with *Xho*I and *Hind*III to form the plasmid pBBlgtB.

### 1.3. Culture conditions

The routine cultures and the preparation of the inocula were performed on LB medium (Sambrook et al., 1989). The high cell density cultures were prepared in a 2-liter fermenter containing an initial volume of 1 liter of medium having the following composition: glycerol (17.5 g.l<sup>-1</sup>) or glucose (15 g.l<sup>-1</sup>), NH<sub>4</sub>H<sub>2</sub>PO<sub>4</sub> (7 g.l<sup>-1</sup>), KH<sub>2</sub>PO<sub>4</sub> (7 g.l<sup>-1</sup>), MgSO<sub>4</sub>·7H<sub>2</sub>O (1 g.l<sup>-1</sup>), thiamine HCl (4.5 mg.l<sup>-1</sup>), solution of trace elements (7.5 ml.l<sup>-1</sup>), citric acid (0.5 g.l<sup>-1</sup>), KOH (2 g.l<sup>-1</sup>). The MgSO<sub>4</sub> is autoclaved separately and the thiamine is sterilized by filtration. The solution of trace elements contains: nitrilotriacetate (70 mM, pH 6.5), ferric citrate (7.5 g.l<sup>-1</sup>), MnCl<sub>2</sub>·4H<sub>2</sub>O (1.3 g.l<sup>-1</sup>), CoCl<sub>2</sub>·6H<sub>2</sub>O (0.21 g.l<sup>-1</sup>), CuCl<sub>2</sub>·2H<sub>2</sub>O (0.13 g.l<sup>-1</sup>), H<sub>3</sub>BO<sub>3</sub> (0.25 g.l<sup>-1</sup>), ZnSO<sub>4</sub>·7H<sub>2</sub>O (1.2 g.l<sup>-1</sup>), Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O (0.15 g.l<sup>-1</sup>). The antibiotics ampicillin (50 mg.l<sup>-1</sup>) and chloramphenicol (25 mg.l<sup>-1</sup>) are added to ensure the presence of the various plasmids. The feed solution contains glycerol (500 g.l<sup>-1</sup>) or glucose (400 g.l<sup>-1</sup>), MgSO<sub>4</sub>·7H<sub>2</sub>O (12 g.l<sup>-1</sup>) and the solution of trace elements (25 ml.l<sup>-1</sup>).

The high cell density cultures are inoculated at 2%. Throughout the culturing, the dissolved oxygen content is maintained at 20% of saturation by manually controlling the flow rate of air and by automatically adjusting the stirring speed. The pH is automatically maintained at 6.8 by addition of aqueous ammonia (15%

w/v). The temperature is maintained at 34°C for the strain JM 109(pCWlgtA) and at 28°C for the strain JM 109(pCWlgtA, pBBlgtB). The high-density culture strategy generally comprises three phases: a first  
5 phase of exponential growth which is ensured by the carbon-based substrate (glycerol or glucose) initially present in the medium; a second phase which starts when the growth becomes limited by the carbon source, which is then added continuously at a rate of 4.5 g.h<sup>-1</sup>.l<sup>-1</sup> of  
10 glycerol or 3.6 g.h<sup>-1</sup>.l<sup>-1</sup> of glucose. In a third phase, this rate is reduced by 60% to slow down the growth, so as to increase the oligosaccharide content.

#### 1.4. Assay of the oligosaccharides

15 Samples (1 ml) are taken during the culturing and are immediately centrifuged in microtubes. The supernatant is retained to assay the extracellular oligosaccharides. The bacterial pellet is resuspended in 1 ml of  
20 water and is then incubated in a water bath at 100°C for 30 minutes to rupture the cells. After a second centrifugation, the supernatant is retained to assay the intracellular oligosaccharides.

25 The lactose concentration is measured using an enzymatic determination kit (Roche diagnostic). The N-acetyl-glucosamine residues present in the oligosaccharides are freed by acid hydrolysis as described previously (Samain et al., 1997) and then  
30 quantified colorimetrically by the method of Reissig et al., (1955); in the description, the term "hydrolyzable GlcNAc" means the amount of GlcNAc assayed in this way.

#### 1.5. Purification of the oligosaccharides

35 At the end of the culturing, the bacterial cells are harvested by centrifugation. The supernatant is retained for purification of the extracellular oligosaccharides. The bacterial cells are resuspended



in 1 liter of water and are then permeabilized by means of a heat treatment (30 minutes at 100°C) to release the intracellular oligosaccharides. After a second centrifugation, these oligosaccharides are recovered in the supernatant.

The first and the second supernatant containing the extracellular and intracellular oligosaccharides, respectively, are adsorbed onto active charcoal (100 g per liter of supernatant). After rinsing with distilled water, the oligosaccharides are diluted with 50% (v/v) ethanol, concentrated by evaporation and freeze-dried.

The oligosaccharides are separated out by steric exclusion chromatography on a column (4.5 cm × 95 cm) of Biogel P4, allowing the injection of about 300 mg of oligosaccharide mixture. The elution is performed with distilled water, at a flow rate of 40 ml.h<sup>-1</sup>.

#### 1.6. Preparation of the allyl β-D-glucosides

Allyl β-D-galactopyranoside and allyl-N-acetyl-β-D-glucosaminide were synthesized according to the protocol described by Lee and Lee (1974).

#### 1.7. Identification and structural characterization of the oligosaccharides

The mass spectra were acquired using a mass spectrometer (Nermag R-1010C). For each experiment, the initial matrix volume is 4 μl. The products were analyzed in FAB<sup>+</sup> mode.

The NMR spectra were obtained using a Bruker AC300 spectrometer.

**Example 2: production of the trisaccharide 4-O-[3-O-(2-acetamido-2-deoxy-β-D-glucopyranosyl)-β-D-**

**galactopyranosyl]-D-glucopyranose, ( $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]-  
 $\beta$ -D-Gal-[1 $\rightarrow$ 4]-D-Glc)**

The principle is illustrated by Figure 1. We used the  
5 strain JM 109 of *Escherichia coli* K12, into which we  
introduced the plasmid pCWlgtA *lgtA* gene. The strain JM  
109 is *lacZ*<sup>-</sup>, that is to say that it is incapable of  
hydrolyzing lactose. On the other hand, it is *lacY*<sup>+</sup>,  
which means that it can synthesize lactose permease.  
10 The *lgtA* gene encodes a  $\beta$ -1,3-N-acetyl-glucosaminyl-  
transferase (LgtA), which transfers an N-acetyl-  
glucosamine unit onto the galactose of lactose.

The strain JM 109 (pCWlgtA) and also the JM 109 control  
15 strain were cultured at high cell density (Samain et  
al., 1997) on glycerol as the carbon and energy  
sources. After the first phase of exponential growth  
provided by the glycerol initially present in the  
medium (17.5 g/l), the growth becomes limited by the  
20 glycerol, which is then added continuously at a rate of  
4.5 g.h<sup>-1</sup>.l<sup>-1</sup>. During this second culturing phase,  
90 mg.h<sup>-1</sup>.l<sup>-1</sup> of lactose are introduced continuously.  
IPTG (isopropyl- $\beta$ -D-thiogalactoside) (0.5 mM) is also  
injected at the start of this phase to induce the  
25 expression of the lactose permease and of the  $\beta$ -1,3-N-  
acetyl-glucosaminyl-transferase. As described in Figure  
2, the added lactose is virtually not accumulated in  
the medium, indicating that the lactose is indeed  
internalized by the bacterial cells. A large  
30 accumulation in the culture medium of a compound  
containing N-acetylglucosamine (hydrolyzable GlcNAc) is  
observed with the strain JM 109 (pCWlgtA). The amount  
of hydrolyzable GlcNAc (3.8 mmol/l) produced  
corresponds almost stoichiometrically to the amount of  
35 lactose consumed (3.5 mmol/l), suggesting that all of  
the lactose internalized has been glycosylated by LgtA.

At the end of the culturing, the cells are removed by  
centrifugation and the oligosaccharides present in the

supernatant are purified by adsorption onto active charcoal and elution with ethanol. The oligosaccharides present are then separated out according to their molecular weight, on a Biogel P4 column. A single  
5 predominant compound is found. The mass spectrometry and NMR data indicate that this compound is indeed the trisaccharide ( $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-[1 $\rightarrow$ 4]- $\beta$ -D-Glc) formed by the addition of a GlcNAc residue to a lactose molecule. Indeed, the mass spectrum in FAB<sup>+</sup> mode shows  
10 the presence of a quasi-molecular ion [M+H]<sup>+</sup> at m/z 546 (Figure 3). The <sup>1</sup>H NMR spectrum confirms the trisaccharide structure, the presence of an acetyl group and the  $\beta$  configuration of the two O-glycoside linkages (Figure 4). The <sup>13</sup>C NMR spectrum also specifies  
15 that the linkage between the GlcNAc and the galactose is indeed of 1,3 type (Figure 5).

### **Example 3: Production of lacto-N-neo-tetraose and of poly lactosamine**

20 The principle is described in Figure 6. The strain of *E. coli* JM 109 was cotransformed with the two plasmids pCWlgtA and pBBlgtB bearing, respectively, the genes *lgtA* (used previously) and *lgtB* (encoding a  $\beta$ -1,4-galactosyl-transferase known as LgtB). The strain JM  
25 109 (pCWlgtA, pBBlgtB) was cultured at high cell density using glucose as the growth substrate. At the start of the second phase, lactose is added at high concentration (5 g.l<sup>-1</sup>) or at low concentration  
30 (1 g.l<sup>-1</sup>) and 0.1 mM IPTG is added. Contrary to what was observed with the strain JM 109 (pCWlgtA), only a weak release of hydrolyzable GlcNAc into the medium is detected during the culturing of this strain. On the other hand, hydrolyzable GlcNAc is found in large  
35 amount in the bacterium (Figure 7). When the supply of lactose is 1 g.l<sup>-1</sup>, complete internalization of the lactose (2.9 mmol.l<sup>-1</sup>) and a total production of bound GlcNAc of 1.45 g.l<sup>-1</sup> (6.5 mmol.l<sup>-1</sup>), i.e. the incorporation of more than 2 GlcNAc molecules per

acceptor lactose molecule, are observed. When the lactose is added in high concentration, the internalization is incomplete ( $3 \text{ g.l}^{-1}$ , i.e.  $8.7 \text{ mmol.l}^{-1}$ ) with a production of GlcNAc also of about 5  $6.5 \text{ mmol.l}^{-1}$ . In this case, the GlcNAc/lactose molar ratio is close to 1, which is coherent with the synthesis of lacto-N-neo-tetraose.

10 The purification of the intracellular oligosaccharide fraction made it possible to obtain several main compounds that are well separated by chromatography on Biogel P4. The mass spectrometry and NMR data indicate that these compounds correspond to the following structures: lacto-N-neo-tetraose  $[M+H]^+ = 708$ ; lacto-N-neo-hexaose  $[M+H]^+ = 708$ ; lacto-N-neo-octaose  $[M+Na]^+ =$  15  $1460$  and probably lacto-N-neo-decaose. The respective proportions of these various compounds depend on the amount of lactose added. Thus, with  $5 \text{ g.l}^{-1}$  of lactose, the major product is lacto-N-neo-tetraose (Figure 8A). 20 On the other hand, a lower supply of lactose ( $1 \text{ g.l}^{-1}$ ) promotes the formation of compounds with a higher degree of polymerization, lacto-N-neo-octaose becoming the major product (Figure 8B).

25 The formation of higher polylactosamine homologs of lacto-N-neo-tetraose is explained by the fact that LgtA is capable of using lacto-N-neo-tetraose to form an intermediate pentasaccharide that is glycosylated by LgtB to give lacto-N-neo-hexaose. The latter compound 30 is itself the precursor for a new glycosylation cycle resulting in the formation of lacto-N-neo-octaose, and so on upto lacto-N-neo-decaose.

35 No significant formation of oligosaccharides with an odd number of residues and bearing a galactose in a nonreducing end position is observed. This indicates that the elongation of the molecules is limited by the incorporation of GlcNAc by LgtA rather than by the galactosylation catalyzed by LgtB.

**Example 4: Production of allyl 3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranoside, ( $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-1 $\rightarrow$ O-allyl)**

5

The strain JM 109(pCWLgtA) was cultured at high cell density on glycerol. At the start of the second phase of culturing, 0.75 g.l<sup>-1</sup> of allyl- $\beta$ -D-galactopyranoside and 0.1 mM of IPTG are added. A total internalization of the allyl- $\beta$ -D-galactopyranoside is observed after 9 hours, with a stoichiometric appearance of hydrolyzable GlcNAc in the extracellular medium. The oligosaccharides present in the extracellular medium are purified as in Example 2. The mass spectrum in FAB<sup>+</sup> mode of the major product obtained shows the presence of a quasi-molecular ion [M+H]<sup>+</sup> at m/z 424 corresponding to the structure  $\beta$ -D-GlcNAc-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-A $\rightarrow$ O-allyl.

20 **Example 5: Production of  $\beta$ -D-Gal-[1 $\rightarrow$ 4]- $\beta$ -D-GlcNAc-1 $\rightarrow$ O-allyl**

The strain JM 109 (pBB1gtB) was cultured at high cell density on glycerol. At the start of the second phase of culturing, 0.5 g.l<sup>-1</sup> of allyl-N-acetyl- $\beta$ -D-glucosaminide ( $\beta$ -D-GlcNAc-1 $\rightarrow$ O-allyl) is added. An approximately 30% reduction in the amount of extracellular hydrolyzable GlcNAc is observed in the first five hours, which demonstrates a partial internalization of allyl-N-acetyl- $\beta$ -D-glucosaminide. In parallel, an almost stoichiometric intracellular production of hydrolyzable GlcNAc and of  $\beta$ -linked galactose residues (hydrolyzable with  $\beta$ -galactosidase) is observed. These results demonstrate that 30% of the allyl-N-acetyl- $\beta$ -D-glucosaminide initially added has been galactosylated by the activity encoded by the *lgtB* gene. After purification, the structure of the expected compound ( $\beta$ -D-Gal-[1 $\rightarrow$ 4]- $\beta$ -D-GlcNAc-1 $\rightarrow$ O-allyl) was confirmed by mass spectrometry and NMR.



REFERENCES

1. Boons (1996) Tetrahedron 52,1095-1121.
- 5 2. Geremia R.A., Mergaert P., Geelen D., Van Montagu M., Holsters M. (1994) Proc. Natl. Acad. Sci. USA 91, 2669-2673.
- 10 3. Gilbert M., Cunningham A.M., Watson D.C., Martin, A., Richards, J.C., Wakarchuk, W.W. (1997) Eur. J. Biochem. 249, 187-194.
- 15 4. Kamst E., van der Drift K.M.G., Thomas-Oates J.E., Lugtenberg B.J.J., Spaink H.P. (1995) Escherichia coli. J. Bacteriol. 177, 6282-6285.
- 20 5. Kovach, M.E., Elzre, P.H., Hill, D.S., Robertson, G.T., Rarris, M.A., Roop II, R.M., Peterson, K.M., (1995). Gene 166, 175-176.
6. Lee R.T., Lee Y.C. (1974). Carbohydr. Res. 37, 193-203.
- 25 7. Martin, S.L., Edbrooke, M.R., Hodgman, T.C., van den Eijnden, D.H., Bird, M.I., (1997) J. Biol. Chem., 34, 21349-21356
- 30 8. Mergaert P., D'Haeze W., Geelen D., Promé D. Van Montagu M., Geremia R., Promé J.C., Holsters M. (1995) J. Biol. Chem. 270, 29217-29223.
9. Reissig, J.L., Strominger, J. L., Leloir, L.F., (1955). J. Biol. Chem. 217, 959-966.
- 35 10. Roy R (1997) Recent developements in the rational design of multivalent glycoconjugates, in Topics curr chem., (eds J.Thiem and H. Driguez), Springer, Heidelberg, pp 241-274.

11. Samain, E., Drouillard, S., Heyraud, A., Driguez, H., Geremia, R.A. (1997). Carbohydr. Res. 30, 235-242.
12. Sambrook, J., Fritsch, E.F., Maniatis, T. (1989)  
5 Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor laboratory Press. N.Y.
13. Spaink H.P., Wijfjes A.H.M., van der Drift K.M.G.,  
Haverkamp J., Thomas-Oates J.E., Lugtenberg B.J.J.  
10 (1994) Mol. Microbiol. 13, 821-831.
14. Yannisch-Perron C., Viera J., Messing J. (1985).  
Gene, 33, 103-119.



## CLAIMS

1. A method for producing an oligosaccharide of interest by a cell starting with at least one  
5 exogenous precursor, said precursor being involved in the biosynthetic pathway of said oligosaccharide, said method comprising the steps of:  
    (i) obtaining a cell that  
10     • comprises at least one gene encoding an enzyme capable of modifying said exogenous precursor or one of the intermediates in the biosynthetic pathway of said oligosaccharide from said exogenous precursor necessary for the synthesis  
15     of said oligosaccharide from said precursor, and also the components for expressing said gene in said cell,  
    • lacks any enzymatic activity liable to degrade said oligosaccharide, said precursor and said  
20     intermediates;  
    (ii) culturing said cell in the presence of at least one said exogenous precursor, under conditions enabling the internalization of said exogenous precursor by said cell and  
25     the production of said oligosaccharide by said cell.
2. The method as claimed in claim 1, characterized in that said cell also comprises at least one gene  
30     encoding an enzyme capable of modifying an endogenous precursor involved in the biosynthetic pathway of said oligosaccharide, said enzyme being identical to or different than the enzyme used in the method described above, and also to the  
35     components for expressing said gene in said cell and characterized in that said cell lacks any enzymatic activity liable to degrade said precursor.

3. The method as claimed in claim 1 or 2, characterized in that said cell is a cell chosen from bacteria and yeasts.
- 5 4. The method as claimed in claim 3, characterized in that the cell is a bacterium, preferably of *Escherichia coli* type.
- 10 5. The method as claimed in one of claims 1 to 4, characterized in that said modification is chosen from glycosylation, sulfatation, acetylation, phosphorylation, succinylation, methylation, and addition of an enolpyruvate group.
- 15 6. The method as claimed in one of claims 1 to 5, characterized in that said enzyme is an enzyme capable of performing a glycosylation, chosen from glycosyl-transferases.
- 20 7. The method as claimed in claim 6, characterized in that said enzyme is a glycosyl-transferase chosen from  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase,  $\beta$ -1,4-galactosyl-transferase,  $\text{l}'\alpha$ -1,3-galactosyl transferase  $\text{l}'\alpha$ -1,4-galactosyl-transferase,  $\text{l}'\alpha$ -2,3-sialyl-transferase,  $\text{l}'\alpha$ -1,3-fucosyl-transferase.
- 25 8. The method as claimed in any one of claims 1 to 7, characterized in that said cell culturing is carried out on a carbon-based substrate.
- 30 9. The method as claimed in claim 8, characterized in that said carbon-based substrate is chosen from glycerol and glucose.
- 35 10. The method as claimed in one of claims 8 and 9, characterized in that said culturing is performed under conditions allowing the production of a culture with a high cell density.

11. The method as claimed in claim 10, characterized in that said culturing step comprises:
- 5 a)- a first phase of exponential cell growth ensured by said carbon-based substrate,
- b)- a second phase of cell growth limited by said carbon-based substrate which is added continuously,
- 10 c)- a third phase of slowed cell growth obtained by continuously adding to the culture an amount of said substrate that is less than the amount of substrate added in step b) so as to increase the content of oligosaccharides produced in the high cell density culture.
- 15
12. The method as claimed in claim 11, characterized in that the amount of substrate added continuously to the cell culture during said phase c) is at least 30% less, preferentially 50% and preferably
- 20 60% less than the amount of substrate added continuously during said phase b).
13. The method as claimed in either of claims 11 and 12, characterized in that said precursor is added
- 25 during phase b).
14. The method as claimed in one of claims 1 to 13, characterized in that said precursor is of carbohydrate nature, preferably of oligosaccharide
- 30 nature.
15. The method according to one of claims 1 to 14 wherein said precursor is internalized by passive transport.
- 35
16. The method as claimed in claim 15, characterized in that said precursor is a monosaccharide whose anomeric carbon is linked to an alkyl group so as

to allow its internalization by a mechanism of passive transport.

5 17. The method as claimed in claim 15, characterized in that said alkyl group is an allyl.

18. The method as claimed in either of claims 15 and 17, for the production of ( $\beta$ -D-Gal-[1 $\rightarrow$ 4]- $\beta$ -D-GlcNac-1 $\rightarrow$ O-allyl), characterized in that  
10 • said cell is a bacterium of *LacZ*<sup>-</sup> genotype;  
• said enzyme is  $\beta$ -1,4-galactosyl-transferase;  
• said substrate is glycerol;  
• said precursor is allyl-N-acetyl- $\beta$ -D-glucosaminide ( $\beta$ -D-GlcNac-1 $\rightarrow$ O-allyl).  
15

19. The method according to one of claims 1 à 14 wherein said precursor is internalized according to active transport.  
20

20. The method as claimed in claim 19, characterized in that said precursor is lactose.

21. The method according to claim 19 wherein said precursor is selected from  
25 • natural or synthetic  $\beta$ -galactosides, such as 4-O- $\beta$ -D-galactopyranosyl-D-fructofuranose (lactulose), 3-O- $\beta$ -D-galacto-pyranosyl-D-arabinose, allyl- $\beta$ -D-galactopyranoside ;  
•  $\alpha$ -galactosides, such as melibiose, raffinose ;  
• saccharose.

30 22. The method as claimed in claims 20 and 21, characterized in that said active transport of said precursor is performed by lactose permease.

35 23. The method as claimed in any one of claims 1 to 22, characterized in that said cell lacks any

enzymatic activity liable to degrade said precursor(s).

- 5           24. The method as claimed in claim 23, characterized in that said cell has a genotype chosen from *LacZ*<sup>-</sup>
- 10           25. The method as claimed in any one of claims 1 to 24, characterized in that it also comprises the addition of an inducer to said culture medium to induce the expression in said cell of said enzyme and/or of a protein involved in said active transport.
- 15           26. The method as claimed in claim 25, characterized in that said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG) and said protein is lactose permease.
- 20           27. The method as claimed in any one of claims 1 to 26, for the production of the trisaccharide 4-O-[3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranosyl]-D-glucopyranose, ( $\beta$ -D-GlcNac-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-[1 $\rightarrow$ 4]-D-Glc), characterized in that:
- 25           • said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype;
- said enzyme is  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase;
- said substrate is glycerol;
- 30           • said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG);
- said precursor is lactose.
- 35           28. The method as claimed in any one of claims 1 to 26, for the production of analogs of lacto-N-neotetraose and of polylactosamines in which the glucose residue is replaced with an allyl group, characterized in that

- said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype;
  - said enzymes are  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase and  $\beta$ -1,4-galactosyl-transferase;
  - 5   • said substrate is glucose;
  - said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG);
  - said precursor is allyl- $\beta$ -D-galactopyranoside.
- 10
29. The method as claimed in any one of claims 1 to 26, for the production of allyl 3-O-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)- $\beta$ -D-galactopyranoside, ( $\beta$ -D-GlcNac-[1 $\rightarrow$ 3]- $\beta$ -D-Gal-1 $\rightarrow$ O-allyl),
- 15 characterized in that:
- said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype;
  - said enzyme is  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase;
  - 20   • said substrate is glycerol;
  - said inducer is isopropyl  $\beta$ -D-thiogalactoside (IPTG);
  - said precursor is allyl- $\beta$ -D-galactopyranoside.
- 25
30. The method as claimed in any one of claims 1 to 26, for the production of lacto-N-neo-tetraose and poly lactosamine (lacto-N-neo-hexaose, lacto-N-neo-octaose, lacto-N-neo-decaose), characterized in
- 30 that:
- said cell is a bacterium of *LacZ*<sup>-</sup>, *LacY*<sup>+</sup> genotype;
  - said enzymes are  $\beta$ -1,3-N-acetyl-glucosaminyl-transferase and  $\beta$ -1,4-galactosyl-transferase;
  - 35   • said substrate is glucose;
  - said inducer is isopropyl- $\beta$ -D-thiogalactoside (IPTG);
  - said precursor is lactose.

31. The method as claimed in claims 1 to 30, for  
5 producing an oligosaccharide labeled with at least  
one isotope, characterized in that said cell is  
cultured on said carbon-based substrate labeled  
with said isotope and/or in the presence of a said  
precursor labeled with said isotope.
- 10 32. An oligosaccharide which may be obtained by the  
method as claimed in any one of claims 1 to 30.
33. An oligosaccharide which may be obtained by the  
method as claimed in any one of claims 18, 21, 29  
15 et 30, characterized in that the double bond of  
the allyl group of said oligosaccharides is  
chemically modified by addition, oxidation or  
ozonolysis reactions to form activated  
oligosaccharides that may be used for the chemical  
20 synthesis of glycoconjugates or glycopolymers.
34. The oligosaccharide as claimed in claim 32 or 33,  
as a medicinal product.
- 25 35. The oligosaccharide as claimed in claim 34, as a  
medicinal product intended to selectively prevent  
the adhesion of biological molecules.
36. The oligosaccharide as claimed in claim 34, as a  
30 medicinal product intended for treating cancer,  
inflammation, heart diseases, diabetes, bacterial  
infections, viral infections and neurological  
diseases and grafts.
- 35 37. A pharmaceutical composition, characterized in  
that it comprises an oligosaccharide as claimed in  
any one of claims 34 to 36 and a pharmaceutically  
acceptable vehicle.

38. The agricultural or agronomic use of an oligosaccharide as claimed in claim 32, especially for the growth and defense of plants.






FIG-1







FIG-2

FIG. 3

FIG.4

FIG. 5




FIG-6






FIG-7

FIG. 8